

ERECTILE DYSFUNCTION TREATMENT (PDE5I)

SILDENAFIL

Viagra®, Vizarsin®, Revatio®

TADALAFIL

Cialis®, Citax®, Adcirca®

VARDENAFIL

Levitra®, Levatik®, Staxyn®

AVANAFIL

Spedra®, Stendra®


	SILDENAFIL	TADALAFIL	VARDENAFIL	AVANAFIL
Dosage	25, 50, 100	2.5, 5 (daily), 10, 20	5, 10, 20	50, 100, 200
Onset	30-60 min	20-30 min	30-40 min	15-20 min FASTEST
Duration	4-5 hours Up to 12 hours	24-36 hours MOST DURABLE	5-6 hours	5-6 hours
Food interaction	Yes HEAVY FATTY MEAL	No	Yes HEAVY FATTY MEAL	Yes HEAVY FATTY MEAL
Side effects	Headache Flushing Abnormal vision	Headache Dyspepsia Back pain/myalgias	Headache Flushing Nasal congestion	Headache Flushing LEAST % SIDE EFFECTS

SILDENAFIL ORAL SUSPENSION (Bandol®, Revatio®):

Bandol® 12.5mg: 25-100mg

50 mg: 4

On demand approx. 1 hour before



Alprostadiil:

-TOPIC-INTRAURETHRAL: MUSE®, VIRIREC®, VITAROS®

ONSET 10 MIN, DURATION 60-90 MIN

-INTRACAVERNOSAL: CAVERJECT®, VIRIDAL®, EDEX®

ONSET 10 MIN, DURATION 40-60 MIN

++PROPOSED CAVERJECT® PROTOCOL:

10---20---30---40---10+15PAPAVERINE---20+30PAPAVERINE

TRIMIX: ALPROSTADIL, PAPAVERINE, PHENTOLAMINE